

Childhood Vaccinations

Information Sheet

- "The greatest threat of childhood dis-eases lies in the dangerous and ineffectual efforts made to prevent them through mass immunizations." *Dr Robert Mendelsohn*

MMIR-Measles, Mumps Rubella:

In Quebec between January and May 1989 a review of 1,600 cases of measles revealed that 58% were vaccinated. *Morbidity & Mortality Weekly Report Vo. 38(A) 1989*

- An Unpublished WHO study on "Measles Susceptible Group" showed that the non-immunized group had a normal contract rate of 2.4%; the immunized group had a contractual level of 33.5% -15 times greater likelihood of infection by the immunized. *Kalokerinos A. and Detti'nan G., Viral Vaccines, Vital or Vulnerable*
- A comprehensive immunization program of Kindergarden to grade 12 students resulted in 61 - 90% occurrence of measles cases in those who received the vaccine. *Maekowitz L. E., New England Journal of Medicine Vol. 320, 1989 pp 75-81*
- A study by the Centre for Disease Control and Prevention in Atlanta revealed a 2.7 times increase in seizure rates within four to seven days the MMR shot and an increase of 3.3 times within 14 days. *What Doctors Won't Tell You, Dec. 1995*

DPT: Diphtheria, Pertussis, Tetanus

In 1975 Japan raised the minimum vaccination **age to 2 years**. This was **followed** by the virtual disappearance of cot death and infant convulsions. Then in the 1980's Japan allowed 3 month and older babies to be vaccinated; the incidence of cot death has increased again. *Vaccination, Viera Scheibner, PhD*

- Whooping cough naturally fell steadily between 1922 - 1975. In 1978 a nationwide immunization program was launched. Almost immediately overall incidence (especially below 6 months) tripled. *Hutchins (1988) Paper presented*
- In 1993 there was a resurgence of pertussis in the United States. Altogether 6,335 cases were reported, the most in 26 years. Since 1993 a pertussis epidemic in Cincinnati occurred primarily among children who had been appropriately immunized, it is clear that the whole-cell pertussis vaccine failed to give full protection against the disease. (*NEngi. JMed. 1994;331:16-21*)

Hepatitis B:

- A large epidemic of diabetes (80% increase) **occurred in New Zealand** following a recent Hepatitis B Immunization program and research scientists believe the most likely explanation is that the immunization program caused the diabetes epidemic. *New Zealand Medical Journal, May 24, 1996 pg. 195.*

Of What Do Vaccine Products Consist?

Specific viruses and bacteria are grown in the following substances, with their foreign proteins (antigens): pig or horse blood, rabbit brain tissue; dog and monkey tissue; chicken and duck egg; and calf serum. N.B. It is generally acknowledged that any foreign substances (including) proteins) which have not been filtered through the body's normal digestive, assimilative and excretory processes, can be highly toxic when freely ranging in the lymphatic and blood systems.

Other foreign additives normally found in various vaccines include:

- * formaldehyde - a known carcinogen
- * thimerosal - an organomercurial antiseptic - 49% mercury
- * aluminum potassium sulfate
- * aluminum phosphate - used in deodorants
- * lactalbumin hydrolysate
- * phenol (carbolic acid) - extremely toxic
- * acetone - volatile and can easily cross the placental barrier
- * glycerin - tri-atomic alcohol derived from decomposed fats which can damage kidney, liver, lungs, local tissue; cause diuresis and possible death

Obomsawin, Raymond, Legitimate Immunity Versus Medical Chaos 1992

R. Moskowitz indicates that "the fact is that we do not know and have never attempted to discover what actually becomes of these foreign substances once they are inside the body." Dr Robert Mendelsohn and Dr Leon Chaitow both have pointed out that long-term effects of immunizations are really unknown.

Alternatives:

Vitamin C

- * Australian microbiologist/physician team of Dettman and Kalokerinos researched the potential of Vitamin C for its efficacy as low cost, safe and wide spectrum anti-viral, anti-toxic and antibacterial agent.
- * Later, North Carolina physician wrote "we have been able to assemble sufficient clinical evidence to prove unequivocally that vitamin C is the antibiotic of choice in the handling of all types of virus diseases."

Vitamin A

- * Scientists at Harvard School of Public Health discovered that vitamin A reduced measles mortality in babies by 90% *Forced Vaccines: Are they here to stay?*, Louise Mollot, *alive magazine* #166p 14
- * Vitamin A treatment of children with measles has proven particularly effective. The WHO now recommends that all children diagnosed with measles in countries with high rates of measles fatalities be given Vitamin A. *Nutrition Today July/August 1992*

Other appropriate alternatives to improve immunity:

- * Physical exercise can enhance natural killer cell function
- * Optimum levels of vitamins, minerals and trace elements
- * A reduction of dietary fat strengthens natural killer cell activity
- * Homeopathy and herbs
- * Adequate sleep improves immunity
- * Reduction of stress levels increase immunity
- * Prudent exposure to ultraviolet rays from sunlight significantly strengthens the immune system

Dr Obomsawin's landmark report, *Universal Immunization... Medical Miracle or Masterful Mirage*

is available through the HANS office for \$14.95 CDN + \$4.50 P&H. This is a well referenced document for anyone interested in the vaccination issue.

- A complete information package on Vaccinations is available from the HANS office for \$20 for HANS members or \$30 for Non-members.

Updated November 20, 1998

Compiled by Health Action Network Society
#202 - 5262 Rumble Street
Burnaby, BC V5J 2B6

(604) 435-0512
1-888-432-4267